

BRITISH COLUMBIA REGIONAL GEOCHEMICAL CLUSTER ANOMALIES AND BEST MATCHES TO MINERAL DEPOSIT TYPES

By C.P. Smyth¹

KEYWORDS: *Geochemistry, Regional Geochemical Survey, RGS, Mineral Deposit Profiles, Mineral Deposit Models, Exploration, Target Generation, Rocks to Riches, MineMatch Geochemistry, MineMatch.*

INTRODUCTION

The “Rocks to Riches” program was inaugurated and funded by the British Columbia government in June 2003 to rekindle mineral investment in the province of British Columbia. It is specifically designed to provide new data or ideas that will attract mineral exploration to BC.

The “MineMatch Geochemistry” project was one of 16 projects approved for funding by the “Rocks to Riches” program in July 2003.

The project’s goal was the generation of Internet-accessible, easy-to-validate exploration targets based on a re-evaluation of all the province’s 45000 regional geochemistry stream sediment sample (RGS) analyses.

All the project’s results have been published on the Internet, and may be freely viewed at www.rockstorichesbc.com.

PROJECT BACKGROUND

Generating high quality exploration targets is a time-consuming and expensive task, particularly when information has to be obtained from multiple sources.

Even in the age of the Internet, integrating this information in a systematic way is a challenge. The MineMatch Geochemistry project seeks to make this integration easier in the context of target generation.

British Columbia’s geology hosts many different mineral deposit types. It also has extensive, diverse, and high quality information records pertinent to minerals exploration and target generation - many of them available on the Internet. The MineMatch Geochemistry project capitalizes on these resources to generate a competitive advantage for companies looking for economic mineral deposits in B.C.

But primarily this project recognizes that, given a supportive permitting and fiscal environment, there can be no better way of encouraging exploration in an area than providing prospectors and companies with sound exploration targets which have not yet been tested. Government geological databases and modern software

techniques can be combined to cost-effectively generate such new, easily validated, exploration targets for free distribution on the Internet to parties interested in exploring in BC.

PROJECT PURPOSE

The purpose of the project was therefore to provide new evidence of potentially economic mineralization in British Columbia in an easy-to-use format, obtained by applying new evaluation methods to the broad coverage of geochemical data existing for the province.

The Internet-accessible maps and supporting documentation delivered by the project are intended to be of immediate value to exploration licence-holders in BC, as well as those seeking to stake new claims in the province.

The project is therefore an example to explorers and potential explorers in BC of how, by investing in a province with an unsurpassed wealth of high-quality, well-maintained base geological data sets, they can gain maximum leverage on their exploration dollars.

PROJECT METHODS

Geochemical Analysis

The project has evaluated the majority of RGS stream sediment sample analyses in the British Columbia Geological Survey (BCGS) RGS database, in conjunction with the sample’s primary-associated rock-types, as derived from the almost-complete integrated 1:250 000 geological map of British Columbia (Massey *et al.*, 2003). Approximately 75% of BC is covered by RGS surveys, as shown in Figure 1. Moss-mat samples were excluded from the project on the basis of their being a different medium from conventional stream sediments.

Geochemical anomaly selection for the study was based on choosing values that exceed the 99th percentile for any specific lithology. Percentile-based threshold selection is widely recognized as the best automated anomaly-picking method in exploration geochemistry (Amor, 2000). A high percentile level was chosen to identify truly anomalous samples.

¹*GeoReference Online Ltd, 301-850 West Hastings St. Vancouver, B.C. V6C 1E1*

TABLE 1
MINEMATCH DESCRIPTION OF ANOMALY CLUSTER NO 3191

Cluster 3191's MineMatch Description	Comment
ElementEnhanced - Au	From Sample 93N831386
ElementEnhanced - Na	From Sample 93N831385
ElementEnhanced - Ta	From Sample 93N831385
ElementEnhanced - U	From Sample 93N831385
RockHost - alkali-feldspar-granite*	From Sample 93N831386
RockHost - granite*	From Sample 93N831386
RockHost - fine-grained-normal-crystalline-rock*	From Sample 93N831385
RockHost - volcaniclastic-igneous-rock*	From Sample 93N831385
* These are the standard rock names taken from the British Geological Survey Rock Classification Scheme (Gillespie <i>et al.</i> , 1999) which most closely match the lithologies shown to be present at the sample sites on the BC 1: 250 000 geological map.	

TABLE 2
SAMPLE DATA FOR CLUSTER NO 3191, WITH SAMPLE POPULATION SIZE AND 99TH PERCENTILE VALUE FOR THE LITHOLOGY TYPES OVER WHICH ITS CONSTITUENT SAMPLES WERE TAKEN

ClusterNo	SampleID	ANOMS	U_INAA	AU1_INAA	NA_INAA	TA_INAA	LITHOLOGY_TYPE
3191	93N831386	1	10	110	2.9	2	granite, alkali feldspar granite intrusive rocks
99th Percentile for Lithology Type		200	94	3.9	20		
No of Samples for Lithology Type		1784	1784	1784	1784		
3191	93N831385	3	18	6	3.7	2.7	undivided volcanic rocks
99th Percentile for Lithology Type		18	164	3.4	2.7		
No of Samples for Lithology Type		1328	1328	1328	1328		

TABLE 3
SIMILARITY RANKING FOR ANOMALY CLUSTER 3191, SHOWING ONLY THE BEST SIX MATCHES

Deposit Type	Similarity Ranking
Subvolcanic Cu-Au-Ag (As-Sb)	1
Porphyry Cu + Mo + Au	2
Hot-spring Au-Ag	3
Porphyry Cu-Au Alkalic	4
Sn Greisen Deposits	5
Gold on flat faults	6
... etcetera	

TABLE 4
EXTRACT FROM THE DETAILED COMPARISON GENERATED BY MINEMATCH OF THE ATTRIBUTES OF ANOMALY CLUSTER 3191 TO THE ATTRIBUTES OF THE BCGS VERSION OF THE PORPHYRY CU/MO/AU DEPOSIT TYPE. THE UNMATCHED DEPOSIT TYPE ATTRIBUTES, MANY OF WHICH DO NOT APPEAR IN THIS EXTRACT, PROVIDE A CHECKLIST OF ATTRIBUTES TO LOOK FOR IN THE VICINITY OF THE ANOMALY CLUSTER

Cluster 3191: Attributes	Cluster 3191's Value	Cluster 3191: Present or Absent	Porphyry Cu + Mo + Au Deposit Type: Attribute	Porphyry Cu + Mo + Au Deposit Type's Value	Deposit Type: Expected Frequency of Attribute Value	Match Type	Porphyry Cu + Mo + Au's Comment
			Alteration	altAssemblage 1	sometimes		Na-Ca silicate alteration, dominated by albite. Sometimes classed as 'propylitic' alteration
ElementEnhanced	Ta	present					
ElementEnhanced	Au	present	ElementEnhanced ToOre	Au	sometimes	maAKOca	Inherited
ElementEnhanced	U	present					
ElementEnhanced	Na	present					
			FormDepositHost	breccia	sometimes		Hydrothermal origin
			FormDepositHost	contact zone	sometimes		Intrusive contact
			FormDepositHost	pluton	usually		Commonly small, multiphase stocks. Deposits referred to as 'classic' or 'southwest U.S.A. type' porphyry coppers
			RockHost	quartz-monzonite	sometimes		Most common intrusive hostrock for mineralization in Canadian Cordillera
RockHost	granite	present	RockHost	granite, porphyritic	sometimes	maybeAKO	
RockHost	fine-grained-normal crystalline-rock	present	RockHost	porphyry	usually	maybeAKO	Commonly granitic plutonic rocks and dykes in intrusive-hydrothermal complex
			RockHost	quartz feldspar porphyry	sometimes		
			RockHost	granodiorite	sometimes		
			RockHost	hornfels	usually		
RockHost	volcaniclastic-igneous-rock	present	RockHost	volcaniclastic-igneous-rock	usually	exact	Commonly andesitic rocks, volcanic flow rocks are locally dominant
RockHost	alkali-feldspar-granite	present	RockHost	<any value>	always	exact	

- (4) Combine lithological information with anomalous cluster characteristics (using spatial joins of sample points with the lithology of the geological polygons within which they lie),

MineMatch[®] was used to:

- (1) Represent information characterizing 95 mineral deposit models;
- (2) Compare anomaly clusters with deposit model descriptions, and publish their similarity rankings; and
- (3) Publish referenced comparison reports (see “Referencing” below) for the best and second-best matching model for each anomaly cluster.

PROJECT OUTPUTS

All project maps, as well as the data used to generate them, are available from the www.rockstorichesbc.com web site.

Geochemical Anomaly Cluster Maps

Figure 2 illustrates the main map output from the project, in which all identified anomaly clusters are shown. These clusters are highly anomalous⁴, as their values exceed 99% of the values for the lithology over which they occur. The clusters are plotted, together with 1:250 000 geology outlines, sample positions, mineral occurrences, and mineral claims boundaries, as they were portrayed on MapPlace in October 2003⁶.

Approximately 85% of the anomaly clusters identified have centroids which lie over free ground, as determined from the mineral claims boundaries mentioned above.

A second way of viewing the project outputs is as maps of anomalous clusters matching different deposit types. Figure 3 shows the distribution of anomalous clusters best matching Copper Porphyry, Eskay Creek Gold, and Zinc-Lead Skarn deposit types.

Reports and References

For each anomaly cluster, the model similarity ranking, best, and second-best match reports are accessed by clicking in the map on the cluster of interest, and then clicking on the desired link in the link-list that appears below the map on the computer screen. See Tables 3 and 4 for examples of these reports.

The best and second-best matching models are hyperlinked to detailed Internet-accessible descriptions of the models on either USGS or BCGS web sites.

Documented mineral occurrences falling within the anomaly cluster boundaries are listed and hyperlinked to their entries in the BCGS MINFILE database.

All anomaly clusters are linked to the MapPlace web site⁵, where custom maps of the samples comprising the anomaly clusters can be dynamically created in a web browser without the need for proprietary software.

Geochemical Sample Statistics

In the course of calculating anomaly thresholds, informative statistical plots were produced, which have value beyond the scope of this project.

For example, bedrock mappers will be interested in the extent to which the statistics validate lumping and splitting of rock types into different mappable lithological units at a scale of 1:250 000. The statistics are also important to environmental studies interested in the background values, and maximum and minimum expected values, of metals in streams in different geological settings. They may also be important to the selection of analytical techniques for future sampling programs.

Consequently all statistical plots produced by the project have been published on the project website, in the “Geochemical Statistics” area.

The statistical plots fall into three categories:

- (a) Box and Whisker plots to summarize compositional distributions for each lithology type;
- (b) Histograms to provide greater detail of compositional distributions for each lithology type; and
- (c) Scatter plots of duplicate analyses.

Figure 4 shows box and whisker plots for cobalt in 31 of the 62 lithologies present on the BC 1:250 000 geology map. These results show how the mean and range of geochemical values change as a function of surrounding lithology-type.

⁴Bearing in mind the qualifications made in the section entitled “Geochemical Analysis” above

⁵A BCGS map portal, accessible at www.em.gov.bc.ca/Mining/Geolsurv/MapPlace/

⁶Some of these claim boundaries might have been up to one year out of date in October 2003. An online titles administration system, scheduled for release in 2004, will remove these backlogs. See the following web site for current titles information www.em.gov.bc.ca/Mining/Titles/TitlesSearch/mguideInfo.htm




Figure 2. Detail from a MineMatch Geochemistry Anomaly Cluster map, showing 2.5 km buffers around anomaly clusters (some may be “clusters” of only one sample), geology polygons, non-anomalous RGS sample points, MINFILE mineral occurrences, and claim outlines. Note that only mineral occurrences within cluster boundaries are referenced in the MineMatch cluster reports.




Figure 3. Distribution of anomaly cluster best-matches, according to deposit type. From left to right: Zn-Pb Skarn, Iron Oxide Copper Gold, and Polymetallic Replacement deposit types.




Figure 4. Box and whisker plots showing the distribution of cobalt levels, as determined by AAS, in 31 of the 62 lithology types appearing on the 1:250000 geology map of British Columbia.




Figure 5. Histograms of chromium levels as determined by Atomic Absorption Spectrometry (left) and Neutron Activation Analysis (right) respectively. All available determinations from the RGS stream sediment database have been plotted. The lower levels in the AA results probably result from only partial extraction of chromium into the solution analysed in the AA spectrometer.

Figure 5 shows histograms of chromium levels as determined by Atomic Absorption Spectrometry and Neutron Activation Analysis respectively. The higher levels in the neutron activation results are almost certainly because of the method's ability to see all the chromium in the sample, while the atomic absorption technique is effectively a partial analysis for chromium.

In addition, scatter plots of all duplicate analyses were generated to assist in anomaly evaluation. A detailed discussion of anomaly evaluation techniques is beyond the scope of this report. However, the importance of these plots may be seen in the differences between the Au (INAA) and V (AAS) duplicates scatter plots in Figures 6a and 6b. Clearly, using the sample medium analysed in the British Columbia RGS program, the absence of an anomalous level of gold in the gold analytical result does not un-equivocally




Figure 6a. Scatter plot of RGS sample duplicate analyses for gold by Neutron Activation.




Figure 6b. Scatter plot of RGS sample duplicate analyses for vanadium by Atomic Absorption Spectrometry.

establish that elevated levels of gold are not present in the sampled stream. This issue has long been recognized in the BCGS, and as a result for many years the BCGS did not analyse for gold because the small samples available were known to be unreliable for gold analysis in many locations because of the nugget effect (Matysek *et al.*, 1988). As with all geochemical surveys, prospectors and geologists should be aware that there can be numerous reasons why an RGS sample can be downstream of a major mineral occurrence, but not show anomalous values.

On the other hand, vanadium analysis of the stream sediments, as measured by AA, which may be reporting only a partial extraction, yields highly reproducible results, as shown in Figure 6b.

CONCLUSION

British Columbia's high quality geological databases and mineral exploration records have been combined with state-of-the-art computer technology to yield a large number of new exploration targets in the province.

These targets have been made available free-of-charge to the world's minerals exploration community, with the purpose of encouraging investment in exploration in British Columbia.

If only one of the new targets leads to an economic discovery, the "Rocks to Riches" program will have paid its sponsors, the taxpayers of British Columbia, a handsome dividend.

ACKNOWLEDGMENTS

Project funding from the Government of British Columbia's "Rocks to Riches" program is gratefully acknowledged.

The author would also like to thank Larry Jones, Nick Massey, Ray Lett and Dave Lefebure, all of the British Columbia Geological Survey, for their assistance in this project.

REFERENCES

Amor, S. (2000): "Data Presentation and Interpretation Requirements for Geochemical Exploration" in Exploration Geochemistry in Today's World: Notes for Professional Development Workshop held at Queen's University, March 11-17, 2000, Part 9, 37 pages.

British Columbia Geological Survey (2001): Five Year Minerals Geoscience Plan for British Columbia, Information Circular 2001-2.

Cox, D. P. and Singer, D. A., Editors (1986): Mineral Deposit Models; U. S. Geological Survey, Bulletin 1693, 379 pages.

- Gillespie, M. R. and Styles, M. T. (1999): BGS Rock Classification Scheme, Volume 1, Classification of Igneous Rocks; British Geological Survey Research Report, (2nd Edition), RR 99-06, 51 pages.
- Lefebure, D. V. and Hoy, T., Editors (1996): Selected British Columbia Mineral Deposit Profiles, Volume 2; B.C. Ministry of Employment and Investment, Open File 1996-13, 172 pages.
- Lefebure, D. V. and Ray, G. E., Editors (1995): Selected British Columbia Mineral Deposit Profiles, Volume 1; B.C. Ministry of Energy, Mines and Petroleum Resources, Open File 1995-20, 135 pages.
- Massey, N.W.D., MacIntyre, D.G. and Desjardins, P.J. (2003): Digital Geology Map of British Columbia: Tile NN10 Central B.C., B.C. Ministry of Energy and Mines, Geofile 2003-21.
- Matysek, P.F. and Day, S.J. (1988): Geochemical Orientation Surveys: Northern Vancouver Island, Fieldwork and preliminary results; B.C. Ministry of Energy, Mines and Petroleum Resources, Geological Fieldwork 1987, Paper 1988-1, pages 493-502.

